


# Gestational Diabetes Screening



## References

1. ACOG Practice Bulletin #137. *Obstet Gynecol* 2013;122:406-16. *Early pregnancy screening for underdiagnosed Type 2 diabetes is suggested for women with certain risk factors.*
2. Lamar ME, Kuehl TJ, Cooney AT. Jelly beans as an alternative to a fifty-gram glucose beverage for gestational diabetes screening. *Am J Obstet Gynecol* 1999; 181:1154-7. *A prospective cohort study of 136 pregnant women showed consistency between one hour glucose values after 28 jelly beans (50g simple carbohydrate) and 50g Glucola.*
3. ACOG Practice Bulletin #137. *Obstet Gynecol* 2013; 122:406-16. *The use of traditional historic factors to identify GDM will miss up to 50%. Low-risk women represent only 10% of the population and selecting these individuals who need not be screened may add unnecessary complexity to the process.*
4. VanDorsten JP et. al. Diagnosing gestational diabetes. NIH Consensus Development Conference Statement 2013;29(1):1-30.
5. Carpenter-Coustan criteria compared with the National Diabetes Data Group thresholds for gestational diabetes; Cheng Y, Block-Kurbisch I; *Obstet Gynecol.* 2009.
6. Bobrowski R, Bottoms S, Micallef J, Dombrowski M. Is the 50-gram glucose screening test ever diagnostic? *J Mat-Fet Med* 1996; 5:317-20. *All subjects with a 50g screen >216 had evidence of GDM and required insulin for glycemic control. Patients with a 50g screen >220 do not all require a 3hr GTT.*
7. Gestational diabetes mellitus. *Diabetes Care* 2004; 27Suppl 1:S88. *A fasting plasma glucose level >126 meets the threshold for diagnosis of diabetes if confirmed on a subsequent day.*
8. Langer O, Anyaegbunam A, Brustman L, Divon M. Management of women with one abnormal oral glucose tolerance test value reduces adverse outcome in pregnancy. *Am J Obstet Gynecol* 1989; 161:593-9. *A RCT including 126 patients with abnormal value on GTT and 146 women in the control group: comparison between the control and the untreated groups showed a significantly higher incidence of large infants and metabolic complications.*

**Revised: November, 2013**

#### Notification to Users

These algorithms are designed to assist the primary care provider in the clinical management of a variety of problems that occur during pregnancy. They should not be interpreted as a standard of care, but instead represent guidelines for management. Variation in practices should take into account such factors as characteristics of the individual patient, health resources, and regional experience with diagnostic and therapeutic modalities.

The algorithms remain the intellectual property of the University of North Carolina at Chapel Hill School of Medicine. They cannot be reproduced in whole or in part without the expressed written permission of the school.